

your guide to emergency contraception

Helping you choose the method
of contraception that is best for you



Emergency contraception

If you have had unprotected sex, that is, sex without using contraception, or think your contraception might have failed, you can use emergency contraception. There are different types of emergency contraception:

- the emergency contraceptive pill, Levonelle
- the emergency contraceptive pill, ellaOne
- the emergency intrauterine device (IUD).

Emergency contraception can be very effective especially if you have an IUD fitted or if the emergency contraceptive pill is taken soon after sex.

You don't need to use emergency contraception for the first 21 days after giving birth.

Q Does emergency contraception cause an abortion?

A

No. Emergency contraception may stop ovulation, fertilisation of an egg, or a fertilised egg from implanting in the uterus (womb). Medical research and legal judgement are quite clear that emergency contraception prevents pregnancy and is not abortion. Abortion can only take place after a fertilised egg has implanted in the uterus.



Where can I get emergency contraception?

You can get emergency contraception free from these places but they may not all supply ellaOne or fit the IUD.

- any general practice that provides contraceptive services
- a contraception clinic
- any young person's service or Brook clinic
- any sexual health clinic
- some genitourinary medicine (GUM) clinics.

You can also get Levonelle free from:

- most NHS walk-in centres (in England only)
- some pharmacies (there may be age restrictions)
- most NHS minor injuries units
- some hospital accident and emergency departments (phone first to check).

You can buy Levonelle from:

- most pharmacies if you are 16 years old or over
- some fee-paying clinics.

Ask your doctor or nurse about getting emergency pills in advance, just in case you need them.

Q How do I buy Levonelle from a pharmacist?

A It will cost around £25. The pharmacist may not be able to sell it to you, for example if:

- it has been more than 72 hours since you had unprotected sex
- you have had unprotected sex more than once in the menstrual cycle
- you think that you might already be pregnant
- you are taking certain prescribed or complementary medicines
- you have certain health conditions.

In these circumstances you will need to see a doctor or nurse. All the advice and treatment you receive is confidential – wherever you receive it.

Q Can someone else get the emergency contraceptive pill for me?

A Someone else will only be given Levonelle on your behalf in exceptional circumstances. You will need to visit the doctor yourself to be prescribed ellaOne.

Q How will I know if my emergency contraception has worked?

A It is unlikely that you will be

pregnant but do a pregnancy test to make sure you are not pregnant if:

- you feel pregnant
- you have not had a normal period within three weeks of taking Levonelle, ellaOne or having the emergency IUD inserted
- you do not have a bleed when you have the seven day break from using the combined pill, contraceptive patch or the contraceptive vaginal ring, or when you take the placebo tablets with EveryDay combined pills.

A pregnancy test will be accurate if the test is done three weeks after the last time you had unprotected sex.

Q Am I protected from future pregnancy?

A

Levonelle and ellaOne will not protect you from pregnancy if you have unprotected sex again. Seek advice – you can take Levonelle as many times as you need to in any menstrual cycle. ellaOne cannot be used more than once in the same menstrual cycle or in the same cycle as taking Levonelle. You can continue to use the IUD as your long-term contraceptive method.

Emergency contraception is not as effective as using other methods of contraception regularly – seek advice on using other methods.



How do I find out about contraception services?

Contact **sexual health direct**, the helpline run by FPA, on 0845 122 8690 or visit www.fpa.org.uk.

- See www.nhs.uk, www.wales.nhs.uk, www.show.scot.nhs.uk or www.n-i.nhs.uk. In England and Wales call NHS Direct on 0845 46 47 and in Scotland NHS 24 on 0845 4 24 24 24.
- In Northern Ireland call the FPA helpline on 0845 122 8687.
- Ask a health centre, pharmacy, hospital, midwife or health visitor.
- Young people can contact Brook on 0808 802 1234, www.brook.org.uk.

Sexually transmitted infections

Most methods of contraception do not protect you from sexually transmitted infections.

Male and female condoms, when used correctly and consistently, can help protect against sexually transmitted infections. If you can, avoid using spermicidally lubricated condoms. The spermicide commonly contains a chemical called Nonoxinol 9 which does not protect against HIV and may even increase the risk of infection.

What is it?

Emergency contraceptive pill – Levonelle

Levonelle is a tablet containing a hormone called progestogen.

You will be given one pill to take. It should be taken within three days (72 hours) of having unprotected sex.

Ask your doctor for advice about taking it within five days (120 hours) of having unprotected sex.

Emergency contraceptive pill – ellaOne

ellaOne is a tablet which contains ulipristal acetate.

You will be given one pill to take within five days (120 hours) of having unprotected sex.

Emergency IUD

An emergency IUD is a small plastic and copper device that is fitted in your uterus up to five days (120 hours) after unprotected sex or within five days of the earliest time you could have released an egg. It takes about 15–20 minutes to fit. It can be uncomfortable – you can ask for a local anaesthetic.

If it cannot be fitted immediately you may be advised to take Levonelle or ellaOne.

Who can use it?

Most women can use Levonelle. However, if you are taking certain prescribed medicines, or complementary medicines you will need advice and the dose of Levonelle may need to be increased. The emergency IUD may be preferred.

Levonelle can be used from day 21 of giving birth. You can use it after a miscarriage or abortion.

Most women can use ellaOne. If you have liver disease, severe asthma or take certain prescribed medicines or complementary medicines, an emergency IUD may be a preferred option.

Avoid breastfeeding for 36 hours after taking ellaOne.

Most women can use an emergency IUD but it is not normally recommended before 28 days after giving birth. If you need to, you can use Levonelle or ellaOne until this time.

You may be able to use the emergency IUD after a miscarriage or abortion. Speak to a doctor or nurse.

What are the disadvantages?

There are no serious short- or long-term side-effects.

- Some women may feel sick, dizzy or tired, or may get headaches, breast tenderness or abdominal pain.
- A very small number will vomit.
- It can disrupt your next period.

Some women may get the same side-effects as for Levonelle above. They may also get:

- painful periods
- mood swings
- muscle and back pain.

Most side-effects go away within a few days.

Some women may get a period-type pain and some light bleeding for a few days after the IUD is fitted. Pain relief can help.

There is a very small chance of getting an infection during the first 20 days after it is fitted. If you already have an infection you may be given antibiotics.

It is not common but the IUD can be pushed out or it can move. There is also a very small risk that it might go through your uterus.

How will it affect my next period?

Your period is likely to come on time or a few days early or late. Sometimes it can be a week late and sometimes even later.

You may have some irregular bleeding between taking Levonelle and your next period. This can range from spotting to being quite heavy.

Your period may be on time, or a few days earlier or later than expected.

Your next period should come at about the same time as you would normally expect it.

Do I need to see a doctor or nurse afterwards?

You should see a doctor or nurse if:

- Your next period is more than seven days late, it is shorter or lighter than usual or you have any sudden or unusual pain in your lower abdomen. These could be signs of an ectopic pregnancy. Although this is not common, it is very serious.
- You are worried that you might have a sexually transmitted infection.

Follow the same advice as for Levonelle above.

You should see a doctor or nurse 3–4 weeks after the IUD is fitted. This is to:

- check you are not pregnant
- discuss your future contraceptive needs
- remove the IUD if this is what you want.

The emergency IUD can be removed during your next period. If removed at any other time you will need to use additional contraception, such as condoms, for seven days before the emergency IUD is taken out.

Can it fail?

Some women get pregnant even though Levonelle was taken correctly. You may also become pregnant if you delay taking it, have further unprotected sex or vomit within two hours of taking it. Speak to your doctor, nurse or pharmacist. They may give you another dose or suggest an emergency IUD.

If you vomit later than two hours Levonelle will have been absorbed.

Some women get pregnant even though they took ellaOne correctly.

You may also become pregnant if you vomit within three hours of taking it. Speak to your doctor or nurse. They may give you another dose or suggest having an emergency IUD. If you vomit later than three hours ellaOne will have been absorbed.

If you cannot feel the IUD threads in the top of your vagina, or you can feel the IUD itself, you may not be protected against pregnancy. See your doctor or nurse straightaway and use additional contraception.

The IUD is very effective but if it fails there is a small increased risk of an ectopic pregnancy. The risk is less in women using an IUD than in women using no contraception at all. Seek advice as soon as possible.

Can I continue to use other contraception?

If you forgot your regular pills or did not use the patch or vaginal ring correctly, you should take your regular pill again, insert a new ring or apply a new patch within 12 hours of taking Levonelle.

Use additional contraception, such as condoms:

- for seven days with the patch, the ring and the combined pill (nine days for Qlaira)
- for two days with the progestogen-only pill.

If you forgot your regular pills or did not use the patch or vaginal ring correctly, you should take your regular pill again, insert a new ring or apply a new patch within 12 hours of taking ellaOne.

Use additional contraception, such as condoms:

- with the patch, the ring and the combined pill for 14 days (16 days for Qlaira)
- with the progestogen-only pill for nine days.

If you want to go back to using your usual contraception, speak to a doctor or nurse about having the IUD removed or you can keep it as your regular method of contraception.

How FPA can help you

Call **sexual health direct**, the helpline run by FPA. It provides:

- confidential information and advice and a wide range of booklets on individual methods of contraception, common sexually transmitted infections, pregnancy choices, abortion and planning a pregnancy
- details of contraception, sexual health and genitourinary medicine (GUM) clinics and sexual assault referral centres.

FPA helplines

England – helpline 0845 122 8690 – 9am to 6pm, Monday to Friday

Northern Ireland – helpline 0845 122 8687 – 9am to 5pm, Monday to Friday

www.fpa.org.uk – visit for confidential information and advice or send your enquiry to Ask WES.

A final word

The information in this booklet is based on evidence-guided research from the World Health Organization and the Faculty of Sexual and Reproductive Healthcare of the Royal College of Obstetricians and Gynaecologists. Different people may give you different information and advice on certain points. All methods of contraception come with a patient information leaflet which provides detailed information about the method.

Remember – contact your doctor, practice nurse or a contraception clinic if you are worried or unsure about anything.



talking sense about sex



Registered charity number 250187. Limited liability company registered in England, number 887632. Supported by the Department of Health.

© FPA printed in March 2011 ISBN: 978-1-905506-98-9